

Integrals**Definitions**

Definite Integral: Suppose $f(x)$ is continuous on $[a, b]$. Divide $[a, b]$ into n subintervals of width Δx and choose x_i^* from each interval.

$$\text{Then } \int_a^b f(x) dx = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i^*) \Delta x.$$

Anti-Derivative: An anti-derivative of $f(x)$ is a function, $F(x)$, such that $F'(x) = f(x)$.

Indefinite Integral: $\int f(x) dx = F(x) + c$ where $F(x)$ is an anti-derivative of $f(x)$.

Fundamental Theorem of Calculus

Part I: If $f(x)$ is continuous on $[a, b]$ then

$$g(x) = \int_a^x f(t) dt \text{ is also continuous on } [a, b]$$

$$\text{and } g'(x) = \frac{d}{dx} \int_a^x f(t) dt = f(x).$$

Part II: $f(x)$ is continuous on $[a, b]$, $F(x)$ is an anti-derivative of $f(x)$ (i.e. $F'(x) = f(x)$)

$$\text{then } \int_a^b f(x) dx = F(b) - F(a).$$

Variants of Part I:

$$\frac{d}{dx} \int_a^{u(x)} f(t) dt = u'(x) f[u(x)]$$

$$\frac{d}{dx} \int_{v(x)}^b f(t) dt = -v'(x) f[v(x)]$$

$$\frac{d}{dx} \int_{v(x)}^{u(x)} f(t) dt = u'(x) f[u(x)] - v'(x) f[v(x)]$$

Properties

$$\int f(x) \pm g(x) dx = \int f(x) dx \pm \int g(x) dx \quad \int cf(x) dx = c \int f(x) dx, c \text{ is a constant}$$

$$\int_a^b f(x) \pm g(x) dx = \int_a^b f(x) dx \pm \int_a^b g(x) dx \quad \int_a^b cf(x) dx = c \int_a^b f(x) dx, c \text{ is a constant}$$

$$\int_a^a f(x) dx = 0 \quad \int_a^b f(x) dx = \int_a^b f(t) dt$$

$$\int_a^b f(x) dx = -\int_b^a f(x) dx \quad \left| \int_a^b f(x) dx \right| \leq \int_a^b |f(x)| dx$$

$$\text{If } f(x) \geq g(x) \text{ on } a \leq x \leq b \text{ then } \int_a^b f(x) dx \geq \int_a^b g(x) dx$$

$$\text{If } f(x) \geq 0 \text{ on } a \leq x \leq b \text{ then } \int_a^b f(x) dx \geq 0$$

$$\text{If } m \leq f(x) \leq M \text{ on } a \leq x \leq b \text{ then } m(b-a) \leq \int_a^b f(x) dx \leq M(b-a)$$

Common Integrals

$$\begin{array}{lll} \int k dx = kx + c & \int \cos u du = \sin u + c & \int \tan u du = \ln |\sec u| + c \\ \int x^n dx = \frac{1}{n+1} x^{n+1} + c, n \neq -1 & \int \sin u du = -\cos u + c & \int \sec u du = \ln |\sec u + \tan u| + c \\ \int x^{-1} dx = \int \frac{1}{x} dx = \ln |x| + c & \int \sec^2 u du = \tan u + c & \int \frac{1}{a^2+u^2} du = \frac{1}{a} \tan^{-1} \left(\frac{u}{a} \right) + c \\ \int \frac{1}{ax+b} dx = \frac{1}{a} \ln |ax+b| + c & \int \sec u \tan u du = \sec u + c & \int \frac{1}{\sqrt{a^2-u^2}} du = \sin^{-1} \left(\frac{u}{a} \right) + c \\ \int \ln u du = u \ln(u) - u + c & \int \csc u \cot u du = -\csc u + c & \\ \int e^u du = e^u + c & \int \csc^2 u du = -\cot u + c & \end{array}$$

Standard Integration Techniques

Note that at many schools all but the Substitution Rule tend to be taught in a Calculus II class.

u Substitution: The substitution $u = g(x)$ will convert $\int_a^b f(g(x))g'(x) dx = \int_{g(a)}^{g(b)} f(u) du$ using $du = g'(x) dx$. For indefinite integrals drop the limits of integration.

$$\begin{array}{ll} \text{Ex. } \int_1^2 5x^2 \cos(x^3) dx & \int_1^2 5x^2 \cos(x^3) dx = \int_1^{\frac{8}{5}} \frac{5}{3} \cos(u) dx \\ u = x^3 \Rightarrow du = 3x^2 dx \Rightarrow x^2 dx = \frac{1}{3} du & = \frac{5}{3} \sin(u) \Big|_1^{\frac{8}{5}} = \frac{5}{3} (\sin(8) - \sin(1)) \\ x=1 \Rightarrow u=1^3=1 \therefore x=2 \Rightarrow u=2^3=8 & \end{array}$$

Integration by Parts: $\int u dv = uv - \int v du$ and $\int_a^b u dv = uv \Big|_a^b - \int_a^b v du$. Choose u and dv from integral and compute du by differentiating u and compute v using $v = \int dv$.

$$\begin{array}{l} \text{Ex. } \int xe^{-x} dx \\ u = x \quad dv = e^{-x} \Rightarrow du = dx \quad v = -e^{-x} \\ \int xe^{-x} dx = -xe^{-x} + \int e^{-x} dx = -xe^{-x} - e^{-x} + c \end{array}$$

$$\begin{array}{l} \text{Ex. } \int_3^5 \ln x dx \\ u = \ln x \quad dv = dx \Rightarrow du = \frac{1}{x} dx \quad v = x \\ \int_3^5 \ln x dx = x \ln x \Big|_3^5 - \int_3^5 \frac{1}{x} dx = (x \ln(x) - x) \Big|_3^5 \\ = 5 \ln(5) - 3 \ln(3) - 2 \end{array}$$

Products and (some) Quotients of Trig Functions

For $\int \sin^n x \cos^m x dx$ we have the following:

- n odd.** Strip 1 sine out and convert rest to cosines using $\sin^2 x = 1 - \cos^2 x$, then use the substitution $u = \cos x$.
- m odd.** Strip 1 cosine out and convert rest to sines using $\cos^2 x = 1 - \sin^2 x$, then use the substitution $u = \sin x$.
- n and m both odd.** Use either 1. or 2.
- n and m both even.** Use double angle and/or half angle formulas to reduce the integral into a form that can be integrated.

Trig Formulas: $\sin(2x) = 2 \sin(x) \cos(x)$, $\cos^2(x) = \frac{1}{2}(1 + \cos(2x))$, $\sin^2(x) = \frac{1}{2}(1 - \cos(2x))$

$$\begin{array}{l} \text{Ex. } \int \tan^3 x \sec^5 x dx \\ \int \tan^3 x \sec^5 x dx = \int \tan^2 x \sec^4 x \tan x \sec x dx \\ = \int (\sec^2 x - 1) \sec^4 x \tan x \sec x dx \\ = \int (u^2 - 1) u^4 du \quad (u = \sec x) \\ = \frac{1}{7} \sec^7 x - \frac{1}{5} \sec^5 x + c \end{array}$$

For $\int \tan^n x \sec^m x dx$ we have the following:

- n odd.** Strip 1 tangent and 1 secant out and convert the rest to secants using $\tan^2 x = \sec^2 x - 1$, then use the substitution $u = \sec x$.
- m even.** Strip 2 secants out and convert rest to tangents using $\sec^2 x = 1 + \tan^2 x$, then use the substitution $u = \tan x$.
- n odd and m even.** Use either 1. or 2.
- n even and m odd.** Each integral will be dealt with differently.

$$\begin{array}{l} \text{Ex. } \int \frac{\sin^5 x}{\cos^3 x} dx \\ \int \frac{\sin^5 x}{\cos^3 x} dx = \int \frac{\sin^4 x \sin x}{\cos^3 x} dx = \int \frac{(\sin^2 x)^2 \sin x}{\cos^3 x} dx \\ = \int \frac{(1 - \cos^2 x)^2 \sin x}{\cos^3 x} dx \quad (u = \cos x) \\ = -\int \frac{(1 - u^2)^2}{u^3} du = -\int \frac{1 - 2u^2 + u^4}{u^3} du \\ = \frac{1}{2} \sec^2 x + 2 \ln |\cos x| - \frac{1}{2} \cos^2 x + c \end{array}$$

Trig Substitutions : If the integral contains the following root use the given substitution and formula to convert into an integral involving trig functions.

$$\sqrt{a^2 - b^2 x^2} \Rightarrow x = \frac{a}{b} \sin \theta \quad \left| \quad \sqrt{b^2 x^2 - a^2} \Rightarrow x = \frac{a}{b} \sec \theta \quad \left| \quad \sqrt{a^2 + b^2 x^2} \Rightarrow x = \frac{a}{b} \tan \theta \right. \right.$$

$$\cos^2 \theta = 1 - \sin^2 \theta \quad \left| \quad \tan^2 \theta = \sec^2 \theta - 1 \quad \left| \quad \sec^2 \theta = 1 + \tan^2 \theta \right. \right.$$

Ex. $\int \frac{16}{x^2 \sqrt{4-9x^2}} dx$

$x = \frac{2}{3} \sin \theta \Rightarrow dx = \frac{2}{3} \cos \theta d\theta$

$\sqrt{4-9x^2} = \sqrt{4-4\sin^2 \theta} = \sqrt{4\cos^2 \theta} = 2|\cos \theta|$

Recall $\sqrt{x^2} = |x|$. Because we have an indefinite integral we'll assume positive and drop absolute value bars. If we had a definite integral we'd need to compute θ 's and remove absolute value bars based on that and,


$$|x| = \begin{cases} x & \text{if } x \geq 0 \\ -x & \text{if } x < 0 \end{cases}$$

In this case we have $\sqrt{4-9x^2} = 2 \cos \theta$.

$\int \frac{16}{x^2 \sqrt{4-9x^2}} dx = \int \frac{16}{\left(\frac{2}{3} \sin \theta\right)^2 (2 \cos \theta)} \left(\frac{2}{3} \cos \theta\right) d\theta = \int \frac{12}{\sin^2 \theta} d\theta$

$= \int 12 \csc^2 \theta d\theta = -12 \cot \theta + c$

Use Right Triangle Trig to go back to x 's. From substitution we have $\sin \theta = \frac{3x}{2}$ so,



From this we see that $\cot \theta = \frac{\sqrt{4-9x^2}}{3x}$. So,

$$\int \frac{16}{x^2 \sqrt{4-9x^2}} dx = -\frac{4\sqrt{4-9x^2}}{x} + c$$

Partial Fractions : If integrating $\int \frac{P(x)}{Q(x)} dx$ where the degree of $P(x)$ is smaller than the degree of $Q(x)$. Factor denominator as completely as possible and find the partial fraction decomposition of the rational expression. Integrate the partial fraction decomposition (P.F.D.). For each factor in the denominator we get term(s) in the decomposition according to the following table.

Factor in $Q(x)$	Term in P.F.D	Factor in $Q(x)$	Term in P.F.D
$ax+b$	$\frac{A}{ax+b}$	$(ax+b)^k$	$\frac{A_1}{ax+b} + \frac{A_2}{(ax+b)^2} + \dots + \frac{A_k}{(ax+b)^k}$
ax^2+bx+c	$\frac{Ax+B}{ax^2+bx+c}$	$(ax^2+bx+c)^k$	$\frac{A_1x+B_1}{ax^2+bx+c} + \dots + \frac{A_kx+B_k}{(ax^2+bx+c)^k}$

Ex. $\int \frac{7x^2+13x}{(x-1)(x^2+4)} dx$

$\int \frac{7x^2+13x}{(x-1)(x^2+4)} dx = \int \frac{4}{x-1} + \frac{3x+16}{x^2+4} dx$

$= \int \frac{4}{x-1} + \frac{3x}{x^2+4} + \frac{16}{x^2+4} dx$

$= 4 \ln|x-1| + \frac{3}{2} \ln(x^2+4) + 8 \tan^{-1}\left(\frac{x}{2}\right)$

Here is partial fraction form and recombined.

$\frac{7x^2+13x}{(x-1)(x^2+4)} = \frac{A}{x-1} + \frac{Bx+C}{x^2+4} = \frac{A(x^2+4)+(Bx+C)(x-1)}{(x-1)(x^2+4)}$

Set numerators equal and collect like terms.

$7x^2+13x = (A+B)x^2 + (C-B)x + 4A - C$

Set coefficients equal to get a system and solve to get constants.


$A+B=7 \quad C-B=13 \quad 4A-C=0$

$A=4 \quad B=3 \quad C=16$

An alternate method that *sometimes* works to find constants. Start with setting numerators equal in previous example : $7x^2+13x = A(x^2+4) + (Bx+C)(x-1)$. Chose *nice* values of x and plug in. For example if $x=1$ we get $20=5A$ which gives $A=4$. This won't always work easily.

Applications of Integrals

Net Area : $\int_a^b f(x) dx$ represents the net area between $f(x)$ and the x -axis with area above x -axis positive and area below x -axis negative.



Area Between Curves : The general formulas for the two main cases for each are,

$y = f(x) \Rightarrow A = \int_a^b [\text{upper function}] - [\text{lower function}] dx$ & $x = f(y) \Rightarrow A = \int_c^d [\text{left function}] - [\text{right function}] dy$

If the curves intersect then the area of each portion must be found individually. Here are some sketches of a couple possible situations and formulas for a couple of possible cases.

$A = \int_a^b f(x) - g(x) dx$ $A = \int_c^d f(y) - g(y) dy$ $A = \int_a^c f(x) - g(x) dx + \int_c^b g(x) - f(x) dx$

Volumes of Revolution : The two main formulas are $V = \int A(x) dx$ and $V = \int A(y) dy$. Here is some general information about each method of computing and some examples.

Rings	Cylinders
$A = \pi((\text{outer radius})^2 - (\text{inner radius})^2)$	$A = 2\pi(\text{radius})(\text{width/height})$
Limits: x/y of right/bot ring to x/y of left/top ring	Limits: x/y of inner cyl. to x/y of outer cyl.
Horz. Axis use $f(x)$, $g(x)$, $A(x)$ and dx .	Horz. Axis use $f(y)$, $g(y)$, $A(y)$ and dy .
Vert. Axis use $f(y)$, $A(y)$ and dy .	Vert. Axis use $f(x)$, $g(x)$, $A(x)$ and dx .

Ex. Axis : $y = a > 0$ **Ex. Axis : $y = a \leq 0$** **Ex. Axis : $y = a > 0$** **Ex. Axis : $y = a \leq 0$**

outer radius : $a - f(x)$ outer radius : $|a| + g(x)$ radius : $a - y$ radius : $|a| + y$

inner radius : $a - g(x)$ inner radius : $|a| + f(x)$ width : $f(y) - g(y)$ width : $f(y) - g(y)$

These are only a few cases for horizontal axis of rotation. If axis of rotation is the x -axis use the $y = a \leq 0$ case with $a = 0$. For vertical axis of rotation ($x = a > 0$ and $x = a \leq 0$) interchange x and y to get appropriate formulas.

Work : If a force of $F(x)$ moves an object in $a \leq x \leq b$, the work done is $W = \int_a^b F(x) dx$

Average Function Value : The average value of $f(x)$ on $a \leq x \leq b$ is $f_{avg} = \frac{1}{b-a} \int_a^b f(x) dx$

Arc Length Surface Area : Note that this is often a Calc II topic. The three basic formulas are,

$$L = \int_a^b ds \quad SA = \int_a^b 2\pi y ds \quad (\text{rotate about } x\text{-axis}) \quad SA = \int_a^b 2\pi x ds \quad (\text{rotate about } y\text{-axis})$$

where ds is dependent upon the form of the function being worked with as follows.

$$ds = \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx \quad \text{if } y = f(x), a \leq x \leq b \quad ds = \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt \quad \text{if } x = f(t), y = g(t), a \leq t \leq b$$

$$ds = \sqrt{1 + \left(\frac{dx}{dy}\right)^2} dy \quad \text{if } x = f(y), a \leq y \leq b \quad ds = \sqrt{r^2 + \left(\frac{dr}{d\theta}\right)^2} d\theta \quad \text{if } r = f(\theta), a \leq \theta \leq b$$

With surface area you *may* have to substitute in for the x or y depending on your choice of ds to match the differential in the ds . With parametric and polar you will always need to substitute.

Improper Integral

An improper integral is an integral with one or more infinite limits and/or discontinuous integrands. Integral is called convergent if the limit exists and has a finite value and divergent if the limit doesn't exist or has infinite value. This is typically a Calc II topic.

Infinite Limit

$$1. \int_a^{\infty} f(x) dx = \lim_{t \rightarrow \infty} \int_a^t f(x) dx \quad 2. \int_{-\infty}^b f(x) dx = \lim_{t \rightarrow -\infty} \int_t^b f(x) dx$$

$$3. \int_{-\infty}^{\infty} f(x) dx = \int_{-\infty}^c f(x) dx + \int_c^{\infty} f(x) dx \quad \text{provided BOTH integrals are convergent.}$$

Discontinuous Integrand

$$1. \text{ Discont. at } a: \int_a^b f(x) dx = \lim_{t \rightarrow a^+} \int_t^b f(x) dx \quad 2. \text{ Discont. at } b: \int_a^b f(x) dx = \lim_{t \rightarrow b^-} \int_a^t f(x) dx$$

$$3. \text{ Discontinuity at } a < c < b: \int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx \quad \text{provided both are convergent.}$$

Comparison Test for Improper Integrals : If $f(x) \geq g(x) \geq 0$ on $[a, \infty)$ then,

$$1. \text{ If } \int_a^{\infty} f(x) dx \text{ conv. then } \int_a^{\infty} g(x) dx \text{ conv.} \quad 2. \text{ If } \int_a^{\infty} g(x) dx \text{ divg. then } \int_a^{\infty} f(x) dx \text{ divg.}$$

Useful fact : If $a > 0$ then $\int_a^{\infty} \frac{1}{x^p} dx$ converges if $p > 1$ and diverges for $p \leq 1$.

Approximating Definite Integrals

For given integral $\int_a^b f(x) dx$ and a n (must be even for Simpson's Rule) define $\Delta x = \frac{b-a}{n}$ and divide $[a, b]$ into n subintervals $[x_0, x_1], [x_1, x_2], \dots, [x_{n-1}, x_n]$ with $x_0 = a$ and $x_n = b$ then,

Midpoint Rule : $\int_a^b f(x) dx \approx \Delta x [f(x_1^*) + f(x_2^*) + \dots + f(x_n^*)]$, x_i^* is midpoint $[x_{i-1}, x_i]$

Trapezoid Rule : $\int_a^b f(x) dx \approx \frac{\Delta x}{2} [f(x_0) + 2f(x_1) + 2f(x_2) + \dots + 2f(x_{n-1}) + f(x_n)]$

Simpson's Rule : $\int_a^b f(x) dx \approx \frac{\Delta x}{3} [f(x_0) + 4f(x_1) + 2f(x_2) + \dots + 2f(x_{n-2}) + 4f(x_{n-1}) + f(x_n)]$